

EVALUATION OF POTATO VARIETIES

and hybrids on the resistance against black scab under conditions of Ukraine's Polissia

Goal. Assess potato varieties and hybrids for resistance against rhizoctonia disease and identify highly resistant varieties. **Methods.** The fundamental principle of the existing methodology for evaluating varieties is to test varieties, hybrids and species of potatoes on natural or artificial infectious backgrounds using a pure culture of the fungus *Rhizoctonia solani* Kuhn. The culture grown in Petri dishes was introduced when the tubers were planted in the soil. The assessment of the degree of damage to tubers and stems was carried out according to the damage to the underground organs of plants, the yield of potatoes, and then the level of variety samples was determined by resistance to the disease. Evaluation of potato varieties and hybrids for resistance against the causative agent of the disease *Rhizoctonia solani* Kuhn was carried out on a nine-point scale, where score 9 — no damage to stems and tubers, 7 — disease development up to 10% (high resistance); 5 — disease progression 11–25% (medium resistance); score 3 — disease progression 26–50% (low resistance). **Results.** Based on the test of potato varieties and hybrids for resistance against rhizoctonia, Dubravka, Vesta, Lugovska, Obriy, Serpanok, Yavir, Scarbnytsia, Tyras and hybrids P.207.407 (Zov × Polisska Rozheva), 205.17-24 (Adretta × 23-16s/73) and others; to medium-resistant (5 points) — Chervona Ruta, Povin, Vodogray, Horlytsia, Zov, Cupava, Nadiyna, Belarossa, Olvia, Svitank Kyivskiyi, Ukrainska Rozheva, Beregynia, Charunka, Okolytsia, Slutch, hybrids P.210.14-23 (Adretta × Posvit), 211.20-31 (Prolisok × Lugovska); to susceptible — 51 or 32.9% of the tested varieties. **Conclusions.** We have proved that mid-early ripening and early ripening cultivars as well as selection hybrids are more infected with black scab than mid-ripening, mid-late ripening and late ripening cultivars. These relatively resistant potato cultivars and hybrids are expedient to use in the selection on the resistance against black scab.

¹V. POLOZHENETS,

Doctor of Agricultural Sciences, professor

²L. NEMERYTSKA,

PhD in Biological Sciences, associated professor

²I. ZHURAVSKA,

PhD in Agricultural Sciences
¹National University of Life and Environmental Sciences of Ukraine, 15, Heroes of Defense str., 03041, Kyiv, Ukraine
²Zhytomyr Agrotechnical Vocational College, 96, Pokrovskaya str., 10031, Zhytomyr, Ukraine

e-mail: ¹familiana@mail.ua,

²luda.nemerizka72@ukr.net,

²innazhuravska1@gmail.com

potato; black scab; disease agent; cultivars; hybrids

Target setting and its connection with important scientific and practical tasks. Black scab is one of the most spread and harmful potato diseases.

It has been proved that fungus *Rhizoctonia solani* Kuhn can infect more than 230 species of plants, but this pathogen can be more often observed on potato [1]. As follows from the researchers' reports black scab of potato was found in England, USA, Russia, Poland, Holland, Bulgaria, Japan and others [2]. In the countries of Western and Eastern Europe, in particular, sclerotium of the agent of *Rhizoctonia solani* Kuhn is observed within 15–30% depending on the resistance degree to the pathogen [3].

According to A.S. Volovyk and colleagues (1995) the potato on the whole territory of Russian Federation is infected with black scab, and the amount of tubers with the symptoms of infection with the agent of *Rhizoctonia solani* Kuhn during the years which are favourable for pathogen development, equals 38% [3].

The results of research in Belarus show that the fungi *Rhizoctonia solani* Kuhn is capable of infec-

ting potato on all stages of ontogenesis [4]. The researchers inform that the agent development depends on weather conditions of a year. Thus, under favourable temperature and humidity the development of fungi on sprouts equals — 25.6–43.3%; on stolons — 23.5–48.2%, on roots — 8.2–27.9% [5].

It has been proven that black scab on potato tubers manifests itself in the form of convex black tubercles, which are sclerotia of the fungus *Rhizoctonia solani* Kuhn, which are easily separated from the periderm of tubers. During the growing season of plants, the disease often manifests itself on the stems in the form of a white stalk, which leads to a violation of the process of photosynthesis, causing the formation of axillary microbulbs of green color [6]. Manifestation of symptoms of black scab on potato tubers is presented in Fig. 1.

The analysis of the latest studies and publications. The latest official publications with the description of potato cultivars contain some information as to their resistance to many diseases, in particular to potato tuber eelworm, potato blight, brown scab, Irish blight, but there is no information as to black scab. The main publications as to potato black scab do not contain the information as to the resistance of most potato cultivars against this disease. The information as to potato resistance against potato blight can not be used for black scab,



Fig. 1. External signs of damage to potato tubers by the fungus *Rhizoctonia solani* Kuhn (Prolisok variety)



because there is some difference between the agents of these diseases [7].

Shaping the goals of the paper — to estimate the resistance of potato cultivars and hybrids against black scab both on the artificial and on natural infection backgrounds under conditions of Ukraine's Polissia as well as to analyze and generalize the results.

Methods. The experiments were conducted at the National University of Life and Environmental Sciences of Ukraine in 2018–2020. Laboratory studies were conducted at the Department of Phytopathology named after Academician V.F. Peresyepkin, and field — on the basis of the research farm «Verzel».

The territory of the experimental field is represented by sod-podzolic gley sandy soil. The particle size distribution of the soil of the experimental field is as follows: 40.5–53.4% sand; 42.4–53.3% dust; 3.7–5.6% mule.

The climate in the experimental field is temperate continental. Perennial amounts of active temperatures are in the range of 2400–2600°C, and the duration of the frost-free period is 150–160 days. The average long-term temperature of the coldest month fluctuates around – 6.4°C. Spring frosts usually end in late April, and the first autumn frosts begin in October. The temperature regime of the summer period is formed under the influence of high pressure coming from the west.

The annual amount of precipitation in the Polissya area of Ukraine is on average 550–650 mm, but in the central part due to the influence of the relief their amount may increase to 900 mm per year. In the cold period (December-March) precipitation is 140–180 mm (20–25% of the annual amount). The lowest amount of precipitation is observed in February and March. The annual amount of precipitation (over the years of research) for the experimental field was 610–680 mm.

Evaluation of potato varieties and breeding hybrids for resistance to black scab was performed on an artificial infectious background using pure culture of the pathogen *Rhizoctonia solani* Kuhn. The fungus of this pathogen was cultured in Petri dishes on potato-glucose medium with the addition of oat extract (1.5%). The five-day-old crop was divided into four parts, and then one of them was

applied to the soil next to the tubers during planting.

Rating of the degree of tuber damage by black scab was performed on a nine-point scale, where at score 9 — no damage to stems and tubers, score 7 — disease development is up to 10% (high resistance); score 5 — disease development 11–25% (average resistance); score 3 — disease development 26–50% (low resistance) [8].

The presentation of the basic material of the research. Most researchers claim that in the purpose of targeted selection of potatoes for resistance to black scab, varieties with extremely high resistance to black scab were not obtained [9–10]. We believe that these circumstances may be due to the broad specialization of the pathogen *Rhizoctonia solani* Kuhn, capable of infecting plants of different families, and therefore it is advisable to select for resistance in the direction of creating varieties with relative resistance to this pathogen.

For field experiments, an artificial infectious background was created by introducing into the soil from Petri dishes an inoculum of the pathogen *Rhizoctonia solani* Kuhn with a load of 8–10 g of pathogen per tuber of the cultivar.

The cultivars of domestic as well as of foreign selection, perspective

selection hybrids as well as some wild, cultural and primitive species (155 cultivar samples in the numbers) were used in testing for estimating of potato cultivars and hybrids resistance against black scab.

From among the tested potato cultivar samples and hybrids 19 or 12.3% belong to relatively resistant ones (7 grades), in particular — Dubravka, Vesta, Lugovska, Obriy, Serpanok, Yavir, Scarbnytsia, Tyras and the hybrids П.207.407 (Polisska Rozheva); 205.17-24 (Adretta × 23-16s/73) and others, 85 cultivar samples or 54.8% belong to mid-resistant (5 grades), in particular — Chervona Ruta, Povin, Vodogray, Horlytsia, Zov, Cupava, Nadiyna, Belarossa, Olivia, Svitanok Kyivskyi, Ukrainska Rozheva, Beregynia, Charunka, Okolytsia, Slutch and the hybrids П.210.14-23 (Adretta × Posvit); 211.20-31 (Prolisok × Lugovska) and 51 or 32.9% belong to susceptible ones (table 1).

The selected cultivar samples of potato with a high resistance degree against black scab are a precious starting material for conducting a goal-oriented selection on the resistance to *Rhizoctonia solani* Kuhn agent (Fig. 1).

The results of the experiments show that the testing of potato varieties for resistance to black scab on an

The list of potato cultivars and hybrids with different degree of resistant to black scab on the artificial infection background (2018–2020)

Cultivar sample	Group of ripening	Degree of tubers infection with black scab		Potential infection, t/ha
		%	бал	
1	2	3	4	5
Relatively resistant				
Dubravka	Mid-ripening	10.4	7.9	40.0–46.0
Lugovska	Mid-ripening	11.3	7.5	47.0–64.0
Obriy	Middle-early	11.5	7.8	45.0–50.0
Serpanok	Early	11.6	7.7	40.0–45.0
Yavir	Mid-ripening	12.0	7.6	40.0–45.0
Scarbnytsia	Early	12.5	7.5	41.0–44.0
Tyras	Early	12.8	7.3	35.0–40.0
П.207.4-7	Middle-early	15.4	7.1	40.0–46.0
П.205.17-24	Mid-ripening	16.0	7.0	42.0–50.0
	<i>HIP₀₅</i>	0,08		1,0–1,1
Mid-resistant				
Chervona Ruta	Middle-late	22.3	6.0	40.0–43.0
Povin	Middle-late	22.4	5.9	41.0–46.0
Vodogray	Mid-ripening	22.7	5.9	40.0–44.0
Gorlytsia	Mid-ripening	22.9	5.7	40.0–46.0
Zov	Early	25.3	5.6	42.0–46.0
Kupava	Middle early	25.9	5.5	41.0–45.0

Continuation of the table.

1	2	3	4	5
Nadiyna	Mid-ripening	26.1	5.4	40.0—46.0
Bellarossa	Mid-ripening	26.4	5.4	45.0—55.0
Svitanok Kyivskyi	Middle-early	26.5	5.5	45.0—50.0
Ukrainska Rozheva	Mid-ripening	27.0	5.4	45.0—50.0
Beregynia	Middle early	28.0	5.3	35.0—45.0
П.210.14.13	Middle early	28.3	5.3	40.0—43.0
П.211.20.31	Middle early	28.6	5.2	34.0—41.5
Okolytsia	Mid-ripening	28.8	5.2	41.0—48.0
Sluch	Mid-ripening	29.0	5.1	45.0—50.0
HIP ₀₅		0,07		0,8—0,9
Susceptive				
Gart	Early	30.6	4.0	35.0—44.0
Posvit	Early	31.4	3.5	33.0—42.0
Prolisok	Early	32.0	3.2	45.0—50.0
Olivia	Mid-late	33.1	3.1	48.5—50.0
Poran	Early	33.9	3.0	41.5—45.0
Ivankivska Rannia	Mid-early	38.0	2.9	40.0—46.0
Slavuta	Under-early	40.2	2.8	39.0—43.0
Kimeria	Early	42.4	2.8	42.0—49.0
HIP ₀₅		0,09		0,7—0,8

artificial infectious background quite objectively characterizes their genetic resistance to the above trait.

CONCLUSIONS

1. It is advisable to evaluate potato varieties and hybrids for resistance to black scab against an artificial infectious background, which makes it possible to objectively determine the genetic resistance against black scab.

2. On the basis of testing of varieties and hybrids of potatoes for resistance to *Rhizoctonia solani* Kuhn, 19 or 12.3% are classified as resistant (7 grades), in particular: — Dubravka, Vesta, Lugovska, Obriy, Serpanok, Yavir, Scarbnysia, Tyras and the hybrids П.207.407 (Polisska Rozheva); 205.17-24 (Adretta × 23-16s/73) and others, 85 cultivar samples or 54.8% belong to mid-resistant (5 grades), in particular — Chervona Ruta, Povin, Vodogray, Horlytsia, Zov, Cupava,

Nadiyna, Belarossa, Olivia, Svitanok Kyivskyi, Ukrainska Rozheva, Beregynia, Charunka, Okolytsia, Sluch and the hybrids П.210.14-23 (Adretta × Posvit); 211.20-31 (Prolisok × Lugovska) and 51 or 32.9% belong to susceptible ones.

Perspectives of a further studies in this direction. A further investigation in this direction lies in a broader studies of other potato cultivars and hybrids as well as in a comparative and complex estimation of the resistance of potato cultivars against black scab which was received by means of other techniques.

REFERENSEC

1. Markov I.L., Bashta O.V., Hentosh D.T., Hlymiaznyi V.A. (2017). Fitopatolohiia : pidruchnyk. [Phytopathology : a textbook]. Kyiv: 548 s. (in Ukrainian).
2. Brehynets O.V. (2015). Kataloh kultur mikroorganizmiv: bakterii, hryby. [Catalog of cultures of microorganisms: bacteria, fungi]. Chernihiv: ISLAV. 47 s. (in Ukrainian).
3. Markov I.L., Ruban M.B. (2014). Dovidnyk iz zakhystu polovykh kultur vid khvorob i shkidnykiv. [Handbook for the protection of field crops from diseases and pests]. Kyiv: Yunivest Media. 384 s. (in Ukrainian).
4. Korniiichuk M.S. (2013). Zakhyst roslyn v adaptovanykh ahrotekhnolohiiakh za optymizatsii zemlekorystuvannia Ukrainy. [Plant protection in adapted agricultural technologies for land use optimization in Ukraine]. Zemlerobstvo. Vyp. 85. S. 100—117. (in Ukrainian).
5. Markov I.L. (2012). Praktykum iz silskohospodarskoi fitopatolohii : navchalnyi posibnyk. [Workshop on agricultural phytopathology: a textbook]. Kyiv: NNTs «Instytut ahrarynoi ekonomiky». 528 s. (in Ukrainian).

6. Nediiska U.I., Semenchuk V.H. (2015). Otsinka produktyvnosti sortiv kartopli. [Estimation of productivity of potato varieties]. Visnyk Tsentru naukovooho zabezpechennia APV Kharkivskoi oblasti. Vyp. 19. S. 143—148. (in Ukrainian).

7. D'yakov Yu.T. (2012). Fundamental'naya fitopatologiya. [Fundamental phytopathology]. Moskva: KRASAND. 512 s. (in Russian).

8. Kutsenko V.S., Osypchuk A.A., Podhaiskyi A.A. et al. 2002. Metodichni rekomendatsii shchodo provedennia doslidzhen z kartopleiu. [Methodical recommendations for conducting research with potatoes]. Nemishaieve: Instytut kartopliarstva UAAN, 183 s. (in Ukrainian).

9. Gentosh D.T., Kyryk M.M., Gentosh I.D., Pikovskiy M.Y. (2020). Species compositions of root rot agents of root rot agents of spring barley. *Ukrainian Journal of Ecology*. 10(3). P. 106—109. doi: 10.15421/2020_141

10. Zakharchuk N.A., Dolya M.M., Tsuman N.V., Stankevych S.V. (2020). Producing of potato varieties resistant to fusarial wilt by cell selection. *Ukrainian Journal of Ecology*. 10(1). P. 289—291. doi: 10.15421/2020_45

¹Положенець В.М., ²Немерицька Л.В., ²Журавська І.А.

¹Національний університет біоресурсів і природокористування України, вул. Героїв Оборони, 15, м. Київ, 03041, Україна

²Житомирський агротехнічний фаховий коледж, вул. Покровська, 96, м. Житомир, 10031, Україна


e-mail: ¹familiana@mail.ua

²luda.nemerizka72@ukr.net,

²innazhuravska1@gmail.com

Оцінювання сортів і гібридів картоплі на стійкість проти ризоктоніозу в умовах Полісся України

Мета. Оцінити сорти й гібриди картоплі на стійкість проти ризоктоніозу та виділити високорезистентні сортозразки. **Методи.** Основний принцип існуючої методології оцінювання сортів зводиться до випробування сортів, гібридів і видів картоплі на природних або штучних інфекційних фонах з використанням чистої культури гриба *Rhizoctonia solani* Kuhn. Вирощену культуру в чашках Петрі вносили при висаджуванні бульб у ґрунт. Бонітування ступеня ураження бульб і стебел здійснювали за ураженням підземних органів рослин, врахувавши картоплі, а далі визначали рівень сортозразків за резистентністю до захворювання. Оцінювали сорти і гібриди картоплі на стійкість проти збудника хвороби *Rhizoctonia solani* Kuhn за дев'ятибальною шкалою: бал 9 — ураження стебел і бульб відсутнє, 7 — розвиток хвороби до 10% (висока стійкість); 5 — розвиток хвороби 11—25% (середня стійкість); бал 3 — розвиток хвороби 26—50% (низька стійкість). **Результати.** На основі випробування сортів і гібридів картоплі на стійкість проти ризоктоніозу до відносно стійких (7 балів) віднесені Дубравка, Веста, Луговська, Обрій, Серпанок, Явір, Скарбніця, Тирас, гібриди П.207.407 (Зов × Поліська рожева), 205.17-24 (Адретта × 23-16с/73) та ін.; до середньостійких (5 балів) — Чervona ruta, Повін, Водограй, Горлиця, Зов, Купава, Надійна, Беллароса, Ольвія, Світанок київський, Українська рожева,



Берегиня, Чарунка, Околиця, Случ, гібриди П.210.14-23 (Адретта × Посвіт), 211.20-31 (Пролисок × Луговська); до сприйнятливих — 51 сорт (32,9% випробовуваних сортів). **Висновки.** Підтверджено, що ранньостиглі і середньоранні сорти та селекційні гібриди значно більше уражуються ризоктоніозом, ніж середньостиглі, середньопізні і пізні. Зазначені відносно стійкі сорти та гібриди картоплі доцільно використовувати у цілеспрямованій селекції на стійкість проти ризоктоніозу.

картопля; ризоктоніоз (чорна парша); збудник хвороби; сорти; гібриди

¹Положенец В.М., ²Немерицкая Л.В.,
³Журавская И.А.

¹Национальный университет биоресурсов и природопользования Украины, ул. Героев Оборона, 15, г. Киев, 03041, Украина

²Житомирский агротехнический профессиональный колледж, ул. Покровская, 96, г. Житомир, 10031, Украина,

e-mail: ¹familiana@mail.ua,

²luda.nemerizka72@ukr.net,

³innazhuravska1@gmail.com

Оценка сортов и гибридов картофеля на устойчивость против ризоктонииоза в условиях Полесья Украины

Цель. Провести оценку сортов и гибридов картофеля на устойчивость против ризоктонииоза и выделить высоко-резистентные сортообразцы. **Методы.** Основопологающий принцип существующей методологии оценивания сортов сводится к испытанию сортов, гибридов и видов картофеля на естественных или искусственных инфекционных фонах с использованием чистой культуры гриба *Rhizoctonia solani* Kühn. Выращенную культуру в чашках Петри вносили при высадке клубней в почву. Бонитировку степени поражения клубней и стеблей осуществляли по поражению подземных органов растений, урожайности картофеля, а затем определяли уровень сортообразцов по резистентности к заболеванию. **Оценку сортов и гибридов картофеля на устойчивость против возбудителя болезни *Rhizoctonia solani* Kühn проводили по девятибалловой шкале, где балл 9 — поражение стеблей и клубней отсутствует, 7 — развитие болезни до 10% (высокая стойкость); 5 — развитие болезни 11—25% (средняя стойкость); балл 3 — развитие болезни 26—50%**

(низкая стойкость). **Результаты.** На основе испытания сортов и гибридов картофеля на устойчивость против ризоктонииоза к относительно устойчивым (7 баллов) отнесены Дубравка, Веста, Луговская, Обрий, Серпанок, Явир, Скарбница, Тирас и гибриды П.207.407 (Зов × Полесская розовая), 205.17-24 (Адретта × 23-16с/73) и др.; к среднеустойчивым (5 баллов) — Червона рута, Повинь, Водограй, Горлиця, Зов, Кунава, Надийна, Беллароса, Ольвия, Свистанок киевский, Украинская розовая, Берегиня, Чарунка, Околиця, Случ, гибриды П.210.14-23 (Адретта × Посвіт), 211.20-31 (Пролисок × Луговская); к восприимчивым — 51, или 32,9% испытываемых сортов. **Выводы.** Подтверждено, что раннеспелые и среднеранние сорта и селекционные гибриды значительно больше поражаются ризоктонииозом, чем среднеспелые, среднепоздние и поздние. Указанные относительно стойкие сорта и гибриды картофеля целесообразно использовать в целеустремленной селекции на стойкость против ризоктонииоза.

картофель; ризоктонииоз (черная парша); возбудитель болезни; сорта; гибриды

Received November 18, 2021